

VYVGART Hytrulo®
(efgartigimod alfa and hyaluronidase-qvfc)

VYVGART®
(efgartigimod alfa-fcab)

VYVGART Dosing Guide

Make the most of a treatment
plan that's designed for you

What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

SELECT IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

DOSING



VYVGART is given in treatment cycles with a personalized break between each cycle


Each treatment cycle is 4 treatments, 1 given each week



Your first treatment cycle



Your personalized break is your time off between receiving treatment cycles



Your next treatment cycle

SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever; have recently received or are scheduled to receive any vaccinations; have any history of allergic reactions; have kidney (renal) problems. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).

HOW PERSONALIZED BREAKS WORK

How personalized breaks may help you get the most out of your treatment plan



You'll work with your doctor to **personalize your breaks** between cycles based on your symptoms and side effects.



When your symptoms start to return, **talk to your doctor** to help them decide when you need another cycle.

For example: Your doctor may start you with a 4-week break after each of your first 3 treatment cycles. Your doctor will continue to adjust your time between additional cycles, based on your symptoms and side effects.

- ➔ The safety of starting the next treatment cycle sooner than 4 weeks from the end of the previous treatment cycle has not been established. If you miss a treatment, VYVGART may be given up to 3 days after the missed treatment. After that, continue with the normal time between treatments.

See what personalized breaks mean for real patients on VYVGART

Visit VYVGARTStories.com to find out more.

SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk. Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to VYVGARTPregnancy.com.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).

SYMPTOM TRACKING



Tracking your symptoms consistently is important

Starting your next treatment cycle is based on your doctor's evaluation of your symptoms and any side effects you may experience. Use the Myasthenia Gravis Activities of Daily Living (MG-ADL) scale to help track your symptoms.



Consistently track your symptoms. If you track them weekly, try to do it at the same time each week.



Use the MG-ADL scale on [page 5](#) to track your symptoms.



Share your MG-ADL score with your doctor to keep them up to date.

Learn more about the importance of symptom tracking at [VYVGARTSymptomTracking.com](https://vvgart.com/symptom-tracking)

SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infection. VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection: fever; chills; frequent and painful urination; cough; pain and blockage of nasal passages; wheezing; shortness of breath; sore throat; excess phlegm; nasal discharge.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).

THE MG-ADL SCALE

The **Myasthenia Gravis Activities of Daily Living (MG-ADL)** scale assesses the impact of generalized myasthenia gravis (gMG) on daily functions by measuring 8 signs or symptoms that are commonly affected in gMG. Each item is measured on a 4-point scale, where a score of 0 represents normal function and a score of 3 represents the loss of ability to perform that function. Total scores range from 0 to 24 points, with a higher score showing more severe gMG.

Grade	0	1	2	3	Score (0, 1, 2, or 3)
1. Talking	Normal	Intermittent slurring or nasal speech	Constant slurring or nasal, but can be understood	Difficult to understand speech	+
2. Chewing	Normal	Fatigue with solid food	Fatigue with soft food	Gastric tube	+
3. Swallowing	Normal	Rare episode of choking	Frequent choking necessitating changes in diet	Gastric tube	+
4. Breathing	Normal	Shortness of breath with exertion	Shortness of breath at rest	Ventilator dependence	+
5. Impairment of ability to brush teeth or comb hair	None	Extra effort, but no rest periods needed	Rest periods needed	Cannot do one of these functions	+
6. Impairment of ability to arise from a chair	None	Mild, sometimes uses arms	Moderate, always uses arms	Severe, requires assistance	+
7. Double vision	None	Occurs, but not daily	Daily, but not constant	Constant	+
8. Eyelid droop	None	Occurs, but not daily	Daily, but not constant	Constant	+
MG-ADL score total = Add lines 1-8					

This information has been provided for educational purposes only and is not meant to be a substitute for professional medical advice. Patients should not use this information for diagnosing a health condition. Patients should always consult a healthcare professional for medical advice or information about diagnosis and treatment.

Source: Wolfe GI, et al. *Neurology*. 1999;52(7):1487-1489.

IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection or fever.
- have recently received or are scheduled to receive any vaccinations.
- have any history of allergic reactions.
- have kidney (renal) problems.
- are pregnant or plan to become pregnant. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby.
- Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to VYVGARTPregnancy.com
- are breastfeeding or plan to breastfeed. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

VYVGART Hytrulo®
(efgartigimod alfa and hyaluronidase-qvfc)

VYVGART®
(efgartigimod alfa-fcab)

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including:

- **Infection.** VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection:
 - fever
 - chills
 - frequent and painful urination
 - cough
 - pain and blockage of nasal passages
 - wheezing
 - shortness of breath
 - sore throat
 - excess phlegm
 - nasal discharge

(Continued on next page)

IMPORTANT SAFETY INFORMATION (cont'd)

VYVGART Hytrulo®
(efgartigimod alfa and hyaluronidase-qvfc)

VYVGART®
(efgartigimod alfa-fcab)

- **Allergic reactions (hypersensitivity reactions).** VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO:

- rash
- swelling of the face, lips, throat, or tongue
- shortness of breath
- trouble breathing
- low blood pressure
- fainting

An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.

- **Infusion or injection-related reactions.** VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction:

- high blood pressure
- chills
- shivering
- chest, stomach, or back pain

The most common side effects of VYVGART or VYVGART HYTRULO include respiratory tract infection, headache, and urinary tract infection.

An additional common side effect with VYVGART HYTRULO includes injection site reactions.

These are not all the possible side effects of VYVGART or VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

Please see full Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection and full Prescribing Information for VYVGART for IV infusion.

Dosage forms and strengths:

VYVGART Hytrulo is available as a single-dose subcutaneous injection containing: 200 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per prefilled syringe, or 180 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per vial. VYVGART is available as a single-dose injection for intravenous use containing 400 mg/20 mL of efgartigimod alfa-fcab per vial.