

VYVGART Hytrulo®
(efgartigimod alfa and hyaluronidase-qvfc)

VYVGART®
(efgartigimod alfa-fcab)

**For MY gMG:
Self-injection with the
VYVGART Hytrulo prefilled
syringe can fit MY plans**

**That's
treatment
MY way**

AChR=acetylcholine receptor.

What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

SELECT IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

#1 prescribed branded biologic treatments for MY gMG*

VYVGART®
(efgartigimod alfa-fcab)

**An IV infusion that's been
FDA-approved for over 3 years**

VYVGART Hytrulo®
(efgartigimod alfa and hyaluronidase-qvfc)

**Self-inject with
the prefilled syringe
at home—or it can
go where you go†**



*VYVGART for IV infusion and VYVGART Hytrulo for subcutaneous injection are the combined #1 prescribed FDA-approved biologic treatments for adults with anti-AChR antibody positive gMG. Based on IQVIA LAAD from January 2023 to December 2024. Data is based on validated open claims of VYVGART for IV infusion, VYVGART Hytrulo for subcutaneous injection, oral products and other biologics that have been approved by the FDA for the treatment of adults with generalized myasthenia gravis. Patients who had more than one medical claim in this dataset were counted only once, based on a pre-defined hierarchy.

†Please see **Patient Information**. Follow appropriate administration steps and storage and handling guidance in the **Instructions for Use**.

SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever; have recently received or are scheduled to receive any vaccinations; have any history of allergic reactions; have kidney (renal) problems. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.



VYVGART can help me do more with MY day 5

Learn about the data from the 2 VYVGART studies



Established safety for VYVGART 8

See safety and side effects from the 2 VYVGART studies



The VYVGART Hytrulo prefilled syringe can fit MY life 10

Learn about the VYVGART Hytrulo prefilled syringe



Things to discuss with MY doctor before starting VYVGART Hytrulo or VYVGART 12

Helpful things to know and talk to your doctor about



A treatment plan that's designed for me 14

Learn about treatment cycles and personalized breaks



My VYVGART Path 16

Support and resources for every step of your journey



Important Safety Information 18

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

Studying the effectiveness and safety of VYVGART



The VYVGART study

- 167 adults received either VYVGART for IV infusion or placebo in addition to their current oral gMG treatment
- Explored changes in daily abilities and muscle weakness as well as safety in adults with anti-AChR antibody positive gMG

The VYVGART Hytrulo study

- 110 adults received either VYVGART Hytrulo for subcutaneous injection or VYVGART for IV infusion in addition to their current oral gMG treatment
- Compared antibody reduction and safety with both treatments in adults with anti-AChR antibody positive gMG

See real patient stories at StoriesOnVYVGART.com

SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk. Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to VYVGARTPregnancy.com.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

THE VYVGART FOR IV INFUSION STUDY

VYVGART can help me do more with MY day

PROVEN TO



IMPROVE DAILY ABILITIES

More than 2x as many patients saw improved daily abilities

68% (44 of 65) of patients who added VYVGART for IV infusion to their current oral gMG treatment significantly improved their ability to perform daily activities during their first treatment cycle, as measured by the Myasthenia Gravis Activities of Daily Living (MG-ADL) scale.* Compared to **30% (19 of 64) of patients on placebo** plus their current oral gMG treatment

PROVEN TO



REDUCE MUSCLE WEAKNESS

More than 4x as many patients saw reduced muscle weakness

63% (41 of 65) of patients who added VYVGART for IV infusion to their current oral gMG treatment saw a significant reduction in muscle weakness during the first treatment cycle, as measured by the Quantitative Myasthenia Gravis (QMG) scale.† Compared to **14% (9 of 64) of patients on placebo** plus their current oral gMG treatment

*Defined as a decrease of 2 or more points on the MG-ADL scale, which measures the severity of 8 common symptoms of gMG, maintained for at least 4 weeks. A lower MG-ADL score means less severe symptoms.

†Defined as a decrease of 3 or more points on the QMG scale, which assesses muscle weakness based on 13 items, maintained for at least 4 weeks. A lower QMG score means less muscle weakness.

SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infection. VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection: fever; chills; frequent and painful urination; cough; pain and blockage of nasal passages; wheezing; shortness of breath; sore throat; excess phlegm; nasal discharge.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

THE VYVGART HYTRULO STUDY

Different delivery, similar effect

Both treatments reduced the harmful AChR antibodies that cause gMG symptoms



VYVGART Hytrulo

reduced harmful
AChR antibodies by

62%



VYVGART

reduced harmful
AChR antibodies by

60%

In the VYVGART for IV infusion study, a reduction in harmful AChR antibodies was associated with improved daily abilities when added to patients' current oral gMG treatment.

VYVGART Hytrulo has 2 active ingredients: efgartigimod alfa, which is the same active ingredient in VYVGART for IV infusion, and hyaluronidase (human recombinant). Hyaluronidase helps to increase the distribution and absorption of ingredients into the body.



SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including allergic reactions (hypersensitivity reactions). VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO: rash; swelling of the face, lips, throat, or tongue; shortness of breath; trouble breathing; low blood pressure; fainting. An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.


Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

THE VYVGART FOR IV INFUSION STUDY

Additional data


The following data point was not the main goal of the study and does not carry the same weight as the main goals of the study. **No reliable conclusions can be drawn about the effectiveness or safety of VYVGART for IV infusion based on this data point.** Talk to your neurologist if you have any questions.

MG-ADL data during the first treatment cycle



of patients who added VYVGART for IV infusion to their current oral gMG treatment **were observed to have an MG-ADL score of 0 or 1, which means minimal or no symptoms**, during at least one visit during their first treatment cycle

vs



of patients on placebo plus their current oral gMG treatment **were observed to have an MG-ADL score of 0 or 1, which means minimal or no symptoms**, during at least one visit during their first treatment cycle


Learn more about the study data for VYVGART Hytrulo for subcutaneous injection and VYVGART for IV infusion at [VYVGARTResults.com](https://www.vyvgartresults.com)

SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infusion or injection-related reactions. VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction: high blood pressure; chills; shivering; chest, stomach, or back pain.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

Established safety for VYVGART for IV infusion



In the VYVGART for IV infusion study, these side effects were reported in **at least 5% of patients on VYVGART** plus their current oral gMG treatment, and more frequently than in patients on placebo plus their current oral gMG treatment.

- Most infections in patients on VYVGART plus their current oral gMG treatment were mild to moderate
- Additionally, more patients on VYVGART compared to placebo plus current oral gMG treatment had lower white blood cell counts that were mild to moderate in severity

Information about allergic, infusion or injection-related reactions

VYVGART or VYVGART Hytrulo can cause allergic reactions such as rash, swelling of the face, lips, throat, or tongue, and shortness of breath. VYVGART Hytrulo can also cause hives. (Continued on page 9.)

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

VYVGART Hytrulo has a similar safety profile



The overall safety of VYVGART Hytrulo for subcutaneous injection was consistent with the safety profile of VYVGART for IV infusion, except patients on VYVGART Hytrulo experienced more injection site reactions.

The safety of VYVGART Hytrulo was established through 2 studies

In the VYVGART for IV infusion study, the most common side effects of VYVGART for IV infusion-treated patients were respiratory tract infection, headache, and urinary tract infection. More patients had lower white blood cell counts that were mild to moderate in severity vs. placebo.

In the VYVGART Hytrulo study, the most common side effects of VYVGART Hytrulo-treated patients were injection site reactions, 38% (21 of 55), and headache. Injection site reactions included injection site rash, redness of the skin, itching sensation, bruising, pain, and hives.

In the VYVGART Hytrulo study and its long-term safety study:

- All injection site reactions were **mild to moderate** and didn't lead to treatment discontinuation
- The majority of injection site reactions **occurred within 24 hours** of injection and resolved on their own
- The majority of injection site reactions **occurred during the first treatment cycle** and decreased with each subsequent cycle

Information about allergic, infusion or injection-related reactions

(Continued from page 8.) VYVGART or VYVGART Hytrulo can also increase the risk of serious allergic reactions such as trouble breathing, low blood pressure, and fainting, as well as infusion or injection-related reactions including high blood pressure, chills, shivering, and chest, stomach, or back pain. These reactions can happen during, shortly after, or weeks after treatment and can cause doctors to pause or, in some cases, stop treatment.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

The VYVGART Hytrulo prefilled syringe can fit MY life, not the other way around



Prefilled and can go where you go

Inject in your own space, whether that's your kitchen table or a hotel room with a view.*

It is injected under your skin (subcutaneously).



Personalized one-on-one training

You and/or your caregiver will receive in-person training at home or in your doctor's office until you're ready to inject on your own.

You'll also get a demonstration kit to help you practice.



Finding more time in your day for the things you love

Self-inject in about 20-30 seconds with the VYVGART Hytrulo prefilled syringe.*

*Please see **Patient Information**. Follow appropriate administration steps and storage and handling guidance in the **Instructions for Use**. Monitor for signs and symptoms of an allergic reaction for at least 30 minutes after injection. If an allergic reaction occurs, you should seek medical attention.

SELECT IMPORTANT SAFETY INFORMATION

The most common side effects of VYVGART or VYVGART HYTRULO include respiratory tract infection, headache, and urinary tract infection. An additional common side effect with VYVGART HYTRULO includes injection site reactions. These are not all the possible side effects of VYVGART or VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.



Amazing
views with my
favorite crew.

That's
treatment
MY way

Learn more about the VYVGART Hytrulo prefilled syringe at [AboutSelfInjections.com](https://www.aboutselfinjections.com)

SELECT IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

Things to discuss with MY doctor before starting VYVGART Hytrulo or VYVGART



No specific vaccines required to start treatment

Tell your doctor if you have recently received or are scheduled to receive any vaccinations.



No routine lab monitoring required

Routine lab monitoring isn't required, but your doctor will still monitor your response to treatment and look out for possible side effects.



Safety

Ask your doctor about the side effects you may experience on VYVGART Hytrulo or VYVGART.



Getting VYVGART Hytrulo your way

With the VYVGART Hytrulo prefilled syringe, you can self-inject, or a caregiver can do it for you. You can still get your injection from a healthcare professional.



Learn about cost and coverage for VYVGART Hytrulo for subcutaneous injection and VYVGART for IV infusion at [VYVGARTAccess.com](https://www.vyvgartaccess.com)

SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever; have recently received or are scheduled to receive any vaccinations; have any history of allergic reactions; have kidney (renal) problems. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

Broad access available for MY treatment*

VYVGART is widely available

90% of commercial and Medicare insured patients have coverage for VYVGART Hytrulo for subcutaneous injection and VYVGART for IV infusion*

Multiple treatment cycles are possible with the first prescription

90% of commercial and Medicare insured patients have coverage allowing for an initial authorization of VYVGART Hytrulo for subcutaneous injection or VYVGART for IV infusion for at least 6 months.* This can allow for multiple treatment cycles with the first prescription

*Based on published policies from Policy Reporter as of August 2024.

SELECT IMPORTANT SAFETY INFORMATION


Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk. Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to VYVGARTPregnancy.com.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

A treatment plan that's designed for me

VYVGART is given in treatment cycles with a personalized break between each cycle

Each treatment cycle is 4 treatments, 1 given each week.




The safety of starting the next treatment cycle sooner than 4 weeks from the end of the previous treatment cycle has not been established. If you miss a treatment, VYVGART may be given up to 3 days after the missed treatment. After that, continue with the normal time between treatments.

SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infection. VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection: fever; chills; frequent and painful urination; cough; pain and blockage of nasal passages; wheezing; shortness of breath; sore throat; excess phlegm; nasal discharge.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

Making the most of personalized breaks



Make the most of a treatment plan that's designed for you. Learn more about treatment cycles and personalized breaks at VYVGARTDosing.com.

Tracking your symptoms consistently is important

Starting your next treatment cycle is based on your doctor's evaluation of your symptoms and any side effects you may experience. Use the MG-ADL scale to help track your symptoms.

Prezina, Kevin, and Doug are paid spokespeople of argenx.

Start tracking with the interactive MG-ADL tool at VYVGARTSymptomTracker.com

SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including allergic reactions (hypersensitivity reactions). VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO: rash; swelling of the face, lips, throat, or tongue; shortness of breath; trouble breathing; low blood pressure; fainting. An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.




Personalized support throughout MY journey

A Nurse Case Manager is here for you

It may feel overwhelming getting started on a treatment. With My VYVGART Path, you'll have access to resources and one-on-one support throughout your treatment journey.


My VYVGART Path is a patient support program that pairs you with a dedicated Nurse Case Manager (NCM), who can help you navigate insurance, offer resources, talk through your challenges, and more.

If you'll be self-injecting with the VYVGART Hytrulo prefilled syringe, an NCM can provide additional education and resources on self-injection.



Information provided by the NCMs is not intended or implied to be a substitute for professional medical advice, diagnosis, or treatment. Always consult with a qualified and licensed physician or other healthcare provider regarding your medical condition, treatment, and healthcare needs.

How MY NCM supports me



A personalized understanding of your treatment

My VYVGART Path NCMs are knowledgeable about VYVGART and are dedicated to listening to you and answering any questions you may have.



Empowering resources and information

NCMs can support you throughout your treatment journey to help you in following your doctor's treatment plan. If you'll be self-injecting with the VYVGART Hytrulo prefilled syringe, they can help with resources for that, too.



Hands-on help navigating the insurance process

Dealing with insurance can be complicated. An NCM can help answer any questions you have, help you understand the process, and verify what is covered by your insurance.



Helpful information about a co-pay program and financial assistance programs

If you have financial concerns or gaps in insurance coverage for your VYVGART therapy, an NCM is here for you with information and support.

Learn about personalized support from an NCM

MyVYVGARTPathgMG.com

1-833-MY-PATH-1 (1-833-697-2841)

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection or fever.
- have recently received or are scheduled to receive any vaccinations.
- have any history of allergic reactions.
- have kidney (renal) problems.
- are pregnant or plan to become pregnant. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby.
 - Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to VYVGARTPregnancy.com
- are breastfeeding or plan to breastfeed. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including:

- **Infection.** VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection,

your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection:

- fever
- chills
- frequent and painful urination
- cough
- pain and blockage of nasal passages
- wheezing
- shortness of breath
- sore throat
- excess phlegm
- nasal discharge

- **Allergic reactions (hypersensitivity reactions).** VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO:

- rash
- swelling of the face, lips, throat, or tongue
- shortness of breath
- trouble breathing
- low blood pressure
- fainting

An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.

- **Infusion or injection-related reactions.** VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction:

- high blood pressure
- chills
- shivering
- chest, stomach, or back pain

The most common side effects of VYVGART or VYVGART HYTRULO include respiratory tract infection, headache, and urinary tract infection. An additional common side effect with VYVGART HYTRULO includes injection site reactions.

These are not all the possible side effects of VYVGART or VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG)

who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

Please see full Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection and full Prescribing Information for VYVGART for IV infusion.

Dosage forms and strengths:

VYVGART Hytrulo is available as a single-dose subcutaneous injection containing: 200 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per prefilled syringe, or 180 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per vial. VYVGART is available as a single-dose injection for intravenous use containing 400 mg/20 mL of efgartigimod alfa-fcab per vial.



Self-injection with the VYVGART Hytrulo prefilled syringe.

That's treatment MY way

- ✓ #1 prescribed branded biologic treatments for adults with anti-AChR antibody positive gMG*
- ✓ VYVGART can improve daily abilities and reduce muscle weakness when added to current oral gMG treatment in adults with anti-AChR antibody positive gMG†
- ✓ VYVGART Hytrulo for subcutaneous injection and VYVGART for IV infusion were safe in treating most patients studied
- ✓ The VYVGART Hytrulo prefilled syringe can go where you go‡

*VYVGART for IV infusion and VYVGART Hytrulo for subcutaneous injection are the combined #1 prescribed FDA-approved biologic treatments for adults with anti-AChR antibody positive gMG. Based on IQVIA LAAD from January 2023 to December 2024. Data is based on validated open claims of VYVGART for IV infusion, VYVGART Hytrulo for subcutaneous injection, oral products and other biologics that have been approved by the FDA for the treatment of adults with generalized myasthenia gravis. Patients who had more than one medical claim in this dataset were counted only once, based on a pre-defined hierarchy.

†Based on the MG-ADL scale, which measures daily abilities, and the QMG scale, which measures muscle weakness.

‡Please see **Patient Information**. Follow appropriate administration steps and storage and handling guidance in the **Instructions for Use**.

SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infusion or injection-related reactions. VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction: high blood pressure; chills; shivering; chest, stomach, or back pain.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

**Ask your
doctor if
VYVGART is
right for you**



**Download the
doctor discussion
guide and start
the conversation**

