

# Challenging gMG with treatment MY way

**VYVGART Hytrulo®**  
(efgartigimod alfa and hyaluronidase-qvfc)

**VYVGART®**  
(efgartigimod alfa-fcab)

FDA-approved for adults  
with any gMG antibody subtype

**Triple**  
seronegative

**Anti-AChR**  
antibody positive

**Anti-LRP4**  
antibody positive

**Anti-MuSK**  
antibody positive

**Monica Seles**  
Tennis legend and member  
of the gMG community

Monica is a paid spokesperson taking VYVGART for intravenous (IV) infusion.

AChR=acetylcholine receptor; LRP4=low-density lipoprotein receptor-related protein 4; MuSK=muscle-specific tyrosine kinase. Triple seronegative means a person does not show anti-AChR, anti-LRP4, or anti-MuSK antibodies when tested.

## What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG).

It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

### SELECT IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.**



# Game on

with the #1 prescribed branded biologic treatments for gMG\*

Based on data for adults with anti-AChR antibody positive gMG

**VYVGART Hytrulo**<sup>®</sup>  
(efgartigimod alfa and hyaluronidase-qvfc)

**VYVGART**<sup>®</sup>  
(efgartigimod alfa-fcab)

\*Combined #1 prescribed FDA-approved branded biologic treatments for adults with anti-AChR antibody positive gMG. Based on IQVIA LAAD from January 2023 to December 2025.

## SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever; have recently received or are scheduled to receive any vaccinations; have any history of allergic reactions; have kidney (renal) problems. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.**

	<b>Studies and proven results</b> .....	<b>4-11</b>
	<b>Safety profiles</b> .....	<b>12-13</b>
	<b>Prefilled syringe for self-injection</b> .....	<b>14</b>
	<b>In-office treatment for IV infusion</b> .....	<b>15</b>
	<b>Dosing</b> .....	<b>16-17</b>
	<b>Things to discuss with your doctor</b> .....	<b>18-19</b>
	<b>My VYVGART Path Patient Support Program</b> .....	<b>20-21</b>
	<b>Important Safety Information</b> .....	<b>22-23</b>

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

## How MY treatment was studied

**A first-of-its-kind treatment for adults with anti-AChR antibody positive gMG**

### ADAPT study



**167 adults received either VYVGART for IV infusion or placebo** in addition to their current oral gMG treatment.



The main goals of the study were to evaluate changes in daily abilities, muscle weakness, and safety.

Learn more about the study at [VYVGARTResults.com](https://www.vyvgartresults.com)

### SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk. Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to [VYVGARTPregnancy.com](https://www.vyvgartpregnancy.com).

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.**

“Small wins are my strongest motivation for staying on VYVGART.”

Monica Seles  
Living with gMC



### SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infection. VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection: fever; chills; frequent and painful urination; cough; pain and blockage of nasal passages; wheezing; shortness of breath; sore throat; excess phlegm; nasal discharge.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).

## I can do more with MY days



### Significantly improved daily abilities\*

- **68% of patients (44 of 65) who added VYVGART for IV infusion** to their current oral gMG treatment significantly improved their daily abilities
- Compared to **30% of patients (19 of 64) who added placebo** to their current oral gMG treatment
- Results measured by the Myasthenia Gravis Activities of Daily Living (MG-ADL) scale in treatment cycle 1 (by week 8)

**That's more than 2x as many patients vs placebo**



\*Defined as a decrease of 2 or more points on the MG-ADL scale, which measures the severity of 8 common symptoms of gMG, maintained for at least 4 weeks. A lower MG-ADL score means less severe symptoms.

### SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including allergic reactions (hypersensitivity reactions). VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO: rash; swelling of the face, lips, throat, or tongue; shortness of breath; trouble breathing; low blood pressure; fainting. An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).**

## Less muscle weakness is a goal



### Significantly reduced muscle weakness†

- **63% of patients (41 of 65) who added VYVGART for IV infusion** to their current oral gMG treatment significantly reduced their muscle weakness
- Compared to **14% of patients (9 of 64) who added placebo** to their current oral gMG treatment
- Results were measured by the Quantitative Myasthenia Gravis (QMG) scale in treatment cycle 1 (by week 8)

**That's more than 4x as many patients vs placebo**



†Defined as a decrease of 3 or more points on the QMG scale, which assesses muscle weakness based on 13 items, maintained for at least 4 weeks. A lower QMG score means less muscle weakness.

### SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infusion or injection-related reactions. VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction: high blood pressure; chills; shivering; chest, stomach, or back pain.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).**

## A milestone moment for MY gMG

The largest study to date supporting FDA approval of VYVGART in adult patients with triple seronegative, anti-LRP4 antibody positive, and anti-MuSK antibody positive gMG

### ADAPT SERON study



**119 adults received either VYVGART for IV infusion or placebo** in addition to their current oral gMG treatment.



The main goals of the study were to compare the difference in daily abilities in patients who received VYVGART for IV infusion vs placebo and safety for the overall patient group.



Actor portrayals inspired by the gMG community.

### SELECT IMPORTANT SAFETY INFORMATION

The most common side effects of VYVGART or VYVGART HYTRULO include respiratory tract infection, headache, and urinary tract infection. An additional common side effect with VYVGART HYTRULO includes injection site reactions. These are not all the possible side effects of VYVGART or VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

## Improvements in daily activities can add up to big victories



### Significantly improved daily abilities

- Patients on VYVGART for IV infusion saw significantly greater improvement in their average MG-ADL score compared to placebo
- **At week 4**, patients on VYVGART for IV infusion saw an average **1.5-point greater improvement** in their daily abilities
- **3.4-point average improvement** of MG-ADL total score for patients who added VYVGART for IV infusion to their current oral gMG treatment compared to a **1.9-point average improvement** of MG-ADL total score for patients who added placebo to their current oral gMG treatment

Learn more about the study at [MoreVYVGARTResults.com](https://www.morevyyvgartresults.com)

### SELECT IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

## A study that gave me choices for MY treatment

**VYVGART Hytrulo was compared to VYVGART for IV infusion in adults with anti-AChR antibody positive gMG**

### ADAPT-SC study



**110 adults received either VYVGART Hytrulo for subcutaneous injection or VYVGART for IV infusion in addition to their current oral gMG treatment.**



The main goals of the study were to compare antibody reduction and safety of both treatments.



### SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever; have recently received or are scheduled to receive any vaccinations; have any history of allergic reactions; have kidney (renal) problems. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.**

## Reducing the main cause of MY gMG symptoms



**VYVGART Hytrulo**  
reduced harmful  
anti-AChR antibodies by

**62%**



**VYVGART for IV infusion**  
reduced harmful  
anti-AChR antibodies by

**60%**

VYVGART Hytrulo and VYVGART for IV infusion were proven to have a similar average reduction in the harmful anti-AChR antibodies that cause gMG symptoms.

### Why antibody reduction was studied


In the **ADAPT study**, a reduction in harmful anti-AChR antibodies was related to improved daily abilities when VYVGART for IV infusion was added to patients' current oral gMG treatment. Improved daily abilities were measured using the MG-ADL scale.

### SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk. Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to [VYVGARTPregnancy.com](http://VYVGARTPregnancy.com).

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.**

## A reliable safety profile for MY treatment



In the **ADAPT study**, these side effects were reported in **at least 5% of patients on VYVGART** plus their current oral gMG treatment and more frequently than in patients on placebo plus their current oral gMG treatment.

- Most infections in patients on VYVGART plus their current oral gMG treatment were mild to moderate
- Additionally, more patients on VYVGART plus current oral gMG treatment compared to placebo plus current oral gMG treatment had lower white blood cell counts that were mild to moderate in severity

### Information about allergic, infusion- or injection-related reactions

VYVGART or VYVGART Hytrulo can cause allergic reactions such as rash, swelling of the face, lips, throat, or tongue, and shortness of breath. VYVGART Hytrulo can also cause hives. *(Continued on page 13.)*

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

## Consistent safety for adults with any gMG antibody subtype



Patients in the **ADAPT SERON study** with triple seronegative, anti-LRP4 antibody positive, or anti-MuSK antibody positive gMG had consistent side effects to those seen in the earlier **ADAPT study** (*page 12*), with the main difference being nausea. 7% of patients on VYVGART plus their current oral gMG treatment experienced nausea compared to 5% of patients on placebo plus their current oral gMG treatment.

## Similar safety for VYVGART Hytrulo and VYVGART for IV infusion



In the **ADAPT-SC study**, the overall safety of VYVGART Hytrulo for subcutaneous injection was similar to the safety profile of VYVGART for IV infusion (*page 12*), except patients on VYVGART Hytrulo plus their current oral gMG treatment experienced more injection site reactions.

In the **ADAPT-SC study**, the most common side effects of VYVGART Hytrulo-treated patients were injection site reactions, 38% (21 of 55), and headache. Injection site reactions included injection site rash, redness of the skin, itching sensation, bruising, pain, and hives.

### Information about allergic, infusion- or injection-related reactions

(Continued from *page 12*.) VYVGART or VYVGART Hytrulo can also increase the risk of serious allergic reactions such as trouble breathing, low blood pressure, and fainting, as well as infusion- or injection-related reactions including high blood pressure, chills, shivering, and chest, stomach, or back pain. These reactions can happen during, shortly after, or weeks after treatment and can cause doctors to pause or, in some cases, stop treatment.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).

# Self-injection can fit MY plans

## VYVGART Hytrulo prefilled syringe for self-injection

“I can self-inject at home or my home away from home.”\*

Ken  
Living with gMG



### Personalized one-on-one training

You and/or your caregiver will receive in-person training at home or in your doctor's office until you're ready to inject on your own. You'll also get a demonstration kit to help you practice.



### More time in your day for the things you love

Self-inject in about 20-30 seconds with the VYVGART Hytrulo prefilled syringe.\* It is injected under your skin (subcutaneously).

## You can choose to get VYVGART Hytrulo for subcutaneous injection from a healthcare professional.

Ken is a paid spokesperson taking VYVGART Hytrulo.

\*Please see **Patient Information**. Follow appropriate administration steps and storage and handling guidance in the **Instructions for Use**. Monitor for signs and symptoms of an allergic reaction for at least 30 minutes after injection. If an allergic reaction occurs, you should seek medical attention.

## SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infection. VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection: fever; chills; frequent and painful urination; cough; pain and blockage of nasal passages; wheezing; shortness of breath; sore throat; excess phlegm; nasal discharge.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

# I can get treatment from MY healthcare team


## VYVGART for IV infusion



Given over 1 hour at an infusion center or doctor's office. In some cases, VYVGART for IV infusion may also be given at home by a trained nurse.†

“After doing my research, I talked to my doctor about VYVGART for IV infusion.”

Monica Seles  
Living with gMG



†A healthcare professional will monitor you for allergic reactions during and for 1 hour after your infusion.

### SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including allergic reactions (hypersensitivity reactions). VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO: rash; swelling of the face, lips, throat, or tongue; shortness of breath; trouble breathing; low blood pressure; fainting. An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).**

## Designed for me and MY gMG symptoms

Each treatment cycle is 4 treatments, 1 given each week.



**You and your doctor will decide together when to schedule your next treatment cycle, taking into account your symptoms and any side effects.**

If you notice a change in your symptoms during your break between cycles, reach out to your doctor to discuss adjusting your dosing schedule. You don't have to wait to start the conversation.

If you miss a treatment, VYVGART may be given up to 3 days after the missed treatment. After that, continue with the normal time between treatments (dosing schedule). Talk to your doctor if you have any questions about your dosing schedule.

### SELECT IMPORTANT SAFETY INFORMATION

VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including infusion or injection-related reactions. VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction: high blood pressure; chills; shivering; chest, stomach, or back pain.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.**

## Making the most of personalized breaks

<p><b>Niki's</b> break between cycles:</p> <p><b>2</b> weeks</p> 	<p><b>Prezina's</b> break between cycles:</p> <p><b>4</b> weeks</p> 	<p><b>Doug's</b> break between cycles:</p> <p><b>9</b> weeks</p> 
<p>On VYVGART for IV infusion since 2022</p>	<p>On VYVGART for IV infusion since 2022</p>	<p>On the VYVGART Hytrulo prefilled syringe since 2025*</p>

### Tracking your symptoms consistently is important

Starting your next treatment cycle is based on your doctor's evaluation of your symptoms and any side effects you may experience. Use the MG-ADL scale to help track your symptoms.

Niki, Prezina, and Doug are paid spokespeople taking VYVGART for IV infusion or VYVGART Hytrulo.

\*Doug started VYVGART for IV infusion in 2022. Doug stopped VYVGART for IV infusion and started self-injection with the VYVGART Hytrulo prefilled syringe in 2025.

Learn more about treatment cycles and personalized breaks at [VYVGARTDosing.com](https://www.vyvgart.com/dosing)

### SELECT IMPORTANT SAFETY INFORMATION

The most common side effects of VYVGART or VYVGART HYTRULO include respiratory tract infection, headache, and urinary tract infection. An additional common side effect with VYVGART HYTRULO includes injection site reactions. These are not all the possible side effects of VYVGART or VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO for subcutaneous injection and full [Prescribing Information](#) for VYVGART for IV infusion.

# Talking to your doctor about starting and staying on VYVGART\*



## No specific vaccines required to start treatment

Tell your doctor if you have recently received or are scheduled to receive any vaccinations before the start of each treatment cycle.



## No regular blood tests required

Regular blood tests aren't required, but your doctor will still monitor your response to treatment and look out for possible side effects.




## Safety

Ask your doctor about the side effects you may experience on VYVGART Hytrulo or VYVGART.



## Getting VYVGART Hytrulo your way

With the VYVGART Hytrulo prefilled syringe, you can self-inject, or a caregiver can do it for you. You can also get your injection from a healthcare professional.



\*For a complete list of what to share with your doctor, refer to the full [Prescribing](#) and [Patient Information](#) for VYVGART Hytrulo, the full [Prescribing Information](#) for VYVGART for IV infusion and the Important Safety Information throughout this brochure.

Download the doctor discussion guide at [VYVGARTGuide.com](https://www.vyvgart.com/guide)

## SELECT IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).

# Broad insurance coverage for VYVGART

## VYVGART is widely available

**>90%** of commercial and Medicare insured patients have coverage for VYVGART Hytrulo for subcutaneous injection and VYVGART for IV infusion.†

## Multiple treatment cycles are possible with the first prescription

**~90%** of commercial and Medicare insured patients have coverage allowing for an initial authorization of VYVGART Hytrulo for subcutaneous injection or VYVGART for IV infusion for at least 6 months.† This can allow for multiple treatment cycles with the first prescription.

†Based on published policies from Policy Reporter as of January 2026 that only included adults with anti-AChR antibody positive gMG.

Learn about cost and coverage at [VYVGARTAccess.com](https://www.vyvgart.com/access)

### SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever; have recently received or are scheduled to receive any vaccinations; have any history of allergic reactions; have kidney (renal) problems. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection](#) and full [Prescribing Information for VYVGART for IV infusion](#).

# My VYVGART® Path

## Personalized support throughout my journey


A Nurse Case Manager (NCM)  
is here for you

It may feel overwhelming getting started on a treatment. With My VYVGART Path, you'll have access to resources and one-on-one support throughout your treatment journey.

**My VYVGART Path is a patient support program** that pairs you with a dedicated NCM. They can help you navigate insurance and potential financial assistance programs, offer resources, talk through your challenges, and assist you in staying on track with your treatment plan.

If you'll be self-injecting with the VYVGART Hytrulo prefilled syringe, an NCM can provide additional education and resources on the process.


Information provided by the NCMs is not intended or implied to be a substitute for professional medical advice, diagnosis, or treatment. Always consult with a qualified and licensed physician or other healthcare provider regarding your medical condition, treatment, and healthcare needs.



“From starting VYVGART and beyond, my Nurse Case Manager guided me at every step.”

Monica Seles  
Living with gMG

## How my Nurse Case Manager supports me



### **A personalized understanding of your treatment**

My VYVGART Path NCMs are knowledgeable about VYVGART and are dedicated to listening to you and answering any questions you may have.



### **Empowering resources and information**

NCMs can support you throughout your treatment journey to help you in following your doctor's treatment plan. If you'll be self-injecting with the VYVGART Hytrulo prefilled syringe, they can help with resources for that, too.



### **Hands-on help navigating the insurance process**

Dealing with insurance can be complicated. An NCM can help answer any questions you have, help you understand the process, and verify what is covered by your insurance.



### **Helpful information about a co-pay program and financial assistance programs**

If you have financial concerns or gaps in insurance coverage for your VYVGART therapy, an NCM is here for you with information and support.

**Learn about personalized support from an NCM**

**[myVYVGARTpath.com](https://myVYVGARTpath.com)**

**[1-833-MY-PATH-1](tel:1-833-MY-PATH-1) (1-833-697-2841)**

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

**Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you:**

- have an infection or fever.
- have recently received or are scheduled to receive any vaccinations.
- have any history of allergic reactions.
- have kidney (renal) problems.
- are pregnant or plan to become pregnant. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby.
  - Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to [VYVGARTPregnancy.com](http://VYVGARTPregnancy.com)
- are breastfeeding or plan to breastfeed. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including:**

- **Infection.** VYVGART or VYVGART HYTRULO may increase the risk of

infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection:

- fever
- chills
- frequent and painful urination
- cough
- pain and blockage of nasal passages
- wheezing
- shortness of breath
- sore throat
- excess phlegm
- nasal discharge

- **Allergic reactions (hypersensitivity reactions).** VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO:

- rash
- swelling of the face, lips, throat, or tongue
- shortness of breath
- trouble breathing
- low blood pressure
- fainting

An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.

- **Infusion or injection-related reactions.** VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction:

- high blood pressure
- chills
- shivering
- chest, stomach, or back pain

**The most common side effects of VYVGART or VYVGART HYTRULO include respiratory tract infection, headache, and urinary tract infection. An additional common side effect with VYVGART HYTRULO includes injection site reactions.**

These are not all the possible side effects of VYVGART or VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?**

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG).

It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

**Please see full Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection and full Prescribing Information for VYVGART for IV infusion.**

### **Dosage forms and strengths:**

VYVGART Hytrulo is available as a single-dose subcutaneous injection containing: 200 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per prefilled syringe, or 180 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per vial. VYVGART is available as a single-dose injection for intravenous use containing 400 mg/20 mL of efgartigimod alfa-fcab per vial.



FDA-APPROVED  
**FOR ADULTS WITH ANY  
gMG ANTIBODY SUBTYPE**

**Triple** seronegative | **Anti-AChR** antibody positive | **Anti-LRP4** antibody positive | **Anti-MuSK** antibody positive

**~5 years prescribed by doctors  
for their patients\***

- VYVGART improved daily abilities when added to patients' current oral gMG treatment†
- A consistent safety profile for any gMG antibody subtype

\*VYVGART for IV infusion has been FDA-approved for adults with anti-AChR antibody positive gMG for ~5 years.

†Based on the MG-ADL scale, which measures daily abilities.

**3 ways to receive treatment**



**Self-injection**  
with the  
VYVGART Hytrulo  
prefilled syringe



**Injection by  
your doctor** with  
VYVGART Hytrulo  
for subcutaneous  
injection



**IV infusion**  
with VYVGART

**SELECT IMPORTANT SAFETY INFORMATION**

Do not use VYVGART® (efgartigimod alfa-fcab) or VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) if you are allergic to any of their ingredients. These medicines may increase infection risk and cause serious allergic, infusion or injection-related reactions.

**Please see Important Safety Information throughout. Please see full Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection and full Prescribing Information for VYVGART for IV infusion.**

VYVGART and VYVGART Hytrulo are registered trademarks of argenx.

For U.S. audiences only.

©2026 argenx US-ESC-22-00011 V6 05/2026

**Ask your  
doctor if  
VYVGART is  
a fit for you**



**Download  
the doctor  
discussion guide**

