

Learn about **self-injection** with the new VYVGART Hytrulo prefilled syringe

Below are answers to commonly asked questions, with more information coming to the website soon.

VYVGART Hytrulo[®]
(efgartigimod alfa and hyaluronidase-qvfc)

→ **What is the VYVGART Hytrulo prefilled syringe?**

VYVGART Hytrulo is a prescription medicine for adults with either chronic inflammatory demyelinating polyneuropathy (CIDP) or anti-AChR antibody positive generalized myasthenia gravis (gMG).

The prefilled syringe contains a single dose of VYVGART Hytrulo. It's now approved for self-injection—which means that VYVGART Hytrulo can be self-injected or injected by a caregiver. It is injected under your skin (subcutaneously).

→ **How do I learn to self-inject?**

You and/or your caregiver will receive in-person training at home or in your doctor's office until you're ready to inject on your own. You'll also get a demonstration kit to help you practice.

→ **Where can I get the VYVGART Hytrulo prefilled syringe?**

After your doctor prescribes VYVGART Hytrulo, your specialty pharmacy will contact you to discuss next steps. You can ask them if delivery to your home or doctor's office is an option for you.*

→ **How long does it take to inject with the VYVGART Hytrulo prefilled syringe?**

It takes about 20-30 seconds to inject with the prefilled syringe. Please see **Patient Information**. Follow appropriate administration steps in the **Instructions for Use**. Monitor for signs and symptoms of an allergic reaction for at least 30 minutes after injection. If an allergic reaction occurs, you should seek medical attention.

→ **Do I always have to be at home to inject VYVGART Hytrulo?**

The prefilled syringe can go where you go. For example, you can inject at your kitchen table or a hotel room with a view.†

→ **What should I ask my doctor if I want to start a conversation about self-injection with the new VYVGART Hytrulo prefilled syringe?**

Talk to your neurologist and ask if self-injection with VYVGART Hytrulo is right for you.

*A specialty pharmacy provides medications that require special handling, storage, and delivery.

†Please see **Patient Information**. Follow appropriate administration steps and storage and handling guidance in the **Instructions for Use**.

AChR=acetylcholine receptor.

SELECT IMPORTANT SAFETY INFORMATION

Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO](#).

IMPORTANT SAFETY INFORMATION

Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Before taking VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection or fever.
- have recently received or are scheduled to receive any vaccinations.
- have any history of allergic reactions.
- have kidney (renal) problems.
- are pregnant or plan to become pregnant. It is not known whether VYVGART HYTRULO will harm your unborn baby.
 - Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to VYVGARTPregnancy.com
- are breastfeeding or plan to breastfeed. It is not known if VYVGART HYTRULO passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

VYVGART HYTRULO can cause side effects which can be serious, including:

- **Infection.** VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection:

- fever	- wheezing
- chills	- shortness of breath
- frequent and painful urination	- sore throat
- cough	- excess phlegm
- pain and blockage of nasal passages	- nasal discharge

• Allergic reactions (hypersensitivity reactions).

VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction:

- | | |
|---|----------------------|
| - rash | - hives |
| - swelling of the face, lips, throat, or tongue | - trouble breathing |
| - shortness of breath | - low blood pressure |
| | - fainting |

• Infusion or injection-related reactions.

VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction:

- | | |
|-----------------------|--------------------------------|
| - high blood pressure | - shivering |
| - chills | - chest, stomach, or back pain |

The most common side effects of VYVGART HYTRULO include respiratory tract infection, headache, urinary tract infection, and injection site reactions.

These are not all the possible side effects of VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

What is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc)?

VYVGART HYTRULO is a prescription medicine used to treat adults with:

- generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.
- chronic inflammatory demyelinating polyneuropathy (CIDP).

It is not known if VYVGART HYTRULO is safe and effective in children.

Please see full Prescribing and Patient Information for VYVGART HYTRULO.

Dosage form and strength:

VYVGART Hytrulo is available as a single-dose subcutaneous injection containing 200 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per prefilled syringe.